

## The Effect of Adding Chinese Food Supplements to Rice on Glycemic Response

Chiao-Hsin Yang, Meng-Hsueh Amanda Lin and Jenshinn Lin\*

Department of Food Science, National Pingtung University of Science and Technology, Neipu, Pingtung 91201, Taiwan (R.O.C.)

\*Corresponding author. E-mail: [jlin@mail.npust.edu.tw](mailto:jlin@mail.npust.edu.tw)

<https://doi.org/10.12982/CMUJNS.2017.0016>

### ABSTRACT

*Although an international table of glycemic index (GI) values has been published, most listed values pertain to single foods. However, in recent years, mixed-diet GI has been widely studied. This study determined the dose-response and mixture interaction effects of food processing supplements (gluten protein, canola oil, and dextrin fiber) combined with white rice on glycemic response and GI. Twelve healthy adults aged 18-26 years were fed a test or reference food containing 50 g of available carbohydrates. Venous blood samples were collected before the meal and at 15, 30, 45, 60, 90, and 120 min after the meal. The results showed that white rice has a GI value of  $93.8 \pm 2.8$ . Supplementation with various amounts of lipids and dextrin reduced the GI slightly, whereas gluten supplementation significantly ( $p < 0.05$ ) reduced the GI from 93.8 to 84.9 and 83.1, respectively. Analysis of the interaction of gluten, lipids, and dextrin on GI using three-way ANOVA revealed that significant effects on GI value were found with gluten ( $p < 0.01$ ) and dextrin ( $p < 0.05$ ). Moreover, adding a gluten protein and dextrin fiber mixture yielded a larger glycemic control effect than adding a lipid and dextrin fiber mixture did. In conclusion, adding gluten protein supplements to rice products may reduce overall postprandial glycemic response and induce a lower GI in healthy people.*

**Keywords:** Glycemic index, White rice, Supplementation, Gluten protein, Dextrin fiber

### INTRODUCTION

Otto and Niklas (1980) were the first to analyze the glycemic responses to various foods. Jenkin et al. (1981) originally conceived the glycemic index (GI). On a scale of 1 to 100, foods can be classified as low GI foods ( $\leq 55$ ), medium GI foods (56 - 69), and high GI foods ( $\geq 70$ ). GIs are typically obtained by dividing the incremental postprandial blood glucose level by the corresponding level after ingesting an equivalent carbohydrate portion (25 or 50 g) of a reference food (Jenkins et al., 1983). Foods with a high GI are more rapidly digested and absorbed, causing greater fluctuations in blood glucose per unit of carbohydrate