

## Effect of a High-Fat Diet and Cholesterol Levels on Depression-like Behavior in Mice

Jutamas Ruanpang<sup>1</sup>, Apisate Pleumsamran<sup>1,2</sup>,  
Juntima Pleumsamran<sup>1,2</sup>, and Salin Mingmalairak<sup>1\*</sup>

<sup>1</sup>Department of Physiology, Faculty of Medicine, Chiang Mai University, Chiang Mai 50200, Thailand

<sup>2</sup>The Northern Neuroscience Center, Maharaj Nakorn Chiang Mai Hospital, Faculty of Medicine, Chiang Mai University, Chiang Mai 50200, Thailand

\*Corresponding author. E-mail: mingmalairak\_s@yahoo.com

<https://doi.org/10.12982/CMUJNS.2018.0012>

### ABSTRACT

*High-fat diets are a well-known risk factor for many diseases, but the correlation between consuming a high-fat diet, serum lipid levels and the severity of depression or depression-like behavior remains unclear. Therefore, this study aimed to determine the effect of a high-fat diet on depression-like behavior and determine the correlation between cholesterol levels and the severity of depressive-like behavior. Thirty-nine, adult, male, C57BL/6Mlac mice were either fed a high-fat (61% of calories from fat) or normal (11% of calories from fat) diet for 10 weeks. At the end of the experiment, serum cholesterol levels and depression-like behavior, as determined by the forced swimming and tail suspension tests, were compared between the mice fed normal (control) and high-fat diets. The mice fed the high-fat diet showed significantly higher total cholesterol, high-density lipoprotein cholesterol (HDL-C), low-density lipoprotein cholesterol (LDL-C), and triglyceride levels, as well as significantly longer immobility times in the forced swimming test (but not the tail suspension test), compared to the control group; thus, we concluded that consuming a high-fat diet for 10 weeks increased serum cholesterol and triglyceride levels and produced depression-like behavior in mice. In the mice fed the high-fat diet, we found large positive correlations between both total cholesterol and HDL-C levels with immobility times during the forced swimming test, indicating a strong link between hyperlipidemia and depression-like behaviors in mice.*

**Keywords:** High-fat diet, Depression-like behavior, Serum lipid level