



Editor:
Wasu Pathom-aree,
Chiang Mai University, Thailand

Article history:
Received: January 28, 2020;
Revised: May 25, 2020;
Accepted: May 27, 2020;
<https://doi.org/10.12982/CMUJNS.2021.002>

Corresponding author:
Tippawan Prapamontol,
e-mail:
tippawan.prapamontol@cmu.ac.th

Research article

Multiple Pesticide Residues Found in Vegetables and Fruits from Rural and Urban Markets in Upper Northern Thailand

Tippawan Prapamontol^{1*}, Surat Hongsibsong¹, Warangkana Naksen^{1,2}, Tanyaporn Kerdnoi¹, Sawaeng Kawichai¹, Watcharapol Polyiem¹, Nisa Pakvilai^{1,3}, Buran Phansawan^{1,4}, and Choochad Santasup⁵

¹ Environment and Health Research Unit, Research Institute for Health Sciences (RIHES), Chiang Mai University, Chiang Mai 50200, Thailand

² Faculty of Public Health, Chiang Mai University, Chiang Mai 50200, Thailand

³ Environmental Science Program, Faculty of Science and Technology, Valaya Alongkorn Rajabhat University, Pathum Thani 13180, Thailand

⁴ Division of Environmental Science, School of Energy and Environment, University of Phayao, Phayao 56000, Thailand

⁵ Department of Plant Science and Natural Resources, Faculty of Agriculture, Chiang Mai University, Chiang Mai 50200, Thailand

Abstract Multiple pesticides are used in vegetable and fruit cultivation worldwide, including in Thailand. A survey series on pesticide residues in vegetable and fruit samples sourced from markets in urban and rural areas of upper northern Thailand during 2007–2013. Sixteen different vegetables (n = 412) and 11 different fruits (n = 301) were analyzed for 43 pesticide residues including 20 organophosphates (OP), 6 synthetic pyrethroids, 12 carbamates, 2 abamectins, imidacloprid, dithiocarbamates, and carbendazim. Out of the 412 vegetable samples, 235 (57%) had pesticide residues and 185 (45%) had pesticide residues that exceeded the maximum residue limits (MRLs). For the fruit samples, 245 (81%) of the 301 samples had pesticide residues and 165 (55%) had pesticide residues that exceeded the MRLs. The vegetable and the fruit samples had multiple synthetic pyrethroid residues and higher levels of residues than OP and other pesticides. Among the OP pesticides, chlorpyrifos was the most frequently detected pesticide. Residue detection in the rural samples was higher than that in the urban samples. The present study found very high numbers of samples to be above the MRLs: 45% of the vegetable samples and 55% of the fruit samples. Therefore, multi-residue methods are proposed as a regular monitoring system to ensure coverage of the multiple pesticides that are commonly used in agriculture and secure the national food safety policy.

Key words; Pesticide residues; Vegetables and fruits; MRL; Food safety

Funding: The present study was part of studies supported by the Thailand Science Research and Innovation (TSRI), formerly the Thailand Research Fund (TRF) (BDG5080018 and RDG5130010) and Chiang Mai University (2007-2010). The authors gratefully acknowledge this support.

Citation: Prapamontol, T., Hongsibsong, S., Naksen, W., Kerdnoi, T., Kawichai, S., Polyiem, W., Pakvilai, N., Phansawan, B., and Santasup, C. 2021. Multiple pesticide residues found in vegetables and fruits from rural and urban markets in upper northern Thailand. *CMUJ. Nat. Sci.* 20(1):e2021002.

INTRODUCTION

Thailand is a tropical country located in Southeast Asia and cultivates several kinds of vegetables and fruits for both domestic consumption and export to places such as Europe (Skretteberg et al., 2015; Poulsen et al., 2017). Multiple pesticides have been reported to be used in vegetable and fruit cultivation in Thailand (Schreinemachers et al., 2011; Schreinemachers and Tipraqsa, 2012; Riwithong et al., 2015) although several sectors have tried to promote reduction of pesticide use (Posri et al., 2006; Chalermphol et al., 2014). The agricultural area in the northern region holds approximately 4 million hectares, or about 22.1%, of the agricultural area of the country. The northern region was ranked the second after the northeastern region (45.4%). In addition, out of the total 5.6 million agricultural farm holdings in the country, the northern region had its farmers using pesticides the most (70%), and ranked second after the central region (73%) (National Statistical Office, 2003). Globally, most of the investigation on multiple pesticide use and residues have been conducted in food sources (Boobis et al., 2008; Prapamontol et al., 2010; Hjorth et al., 2011; Jardim and Caldas, 2012; Nougadère et al., 2012; Arias et al., 2014; Panseri et al., 2014; Szpyrka et al., 2015; Wanwimolruk et al., 2019), and some have been carried out in environmental samples (Sangchan et al., 2012; Pose-Juan et al., 2015). Biological assessment of exposure to pesticides has been reported from various populations in northern Thailand such as small-scale farmers (Panuwet et al., 2008; Hongsibsong et al., 2019), school children (Panuwet et al., 2009), farmers and consumers (Thiphom et al., 2014; Wongta et al., 2018), and nursing farmers (Naksen et al., 2014). Urinary pesticide metabolites found among small-scale farmers were metabolites of organophosphate (OP) pesticides, for example, dialkylphosphates (DAPs) (Hongsibsong et al., 2019); ethylene bisdithiocarbamates, e.g., ethylenethiourea (ETU); malathion; 2,4-D; and alachlor (Panuwet et al., 2008). Whereas the specific urinary metabolites detected in school children for example para-nitrophenol (PNP); 3,5,6-trichloro-2-pyridinol (TPCY, metabolite of chlorpyrifos); and 3-phenoxybenzoic acid (3-PBA, a common metabolite of pyrethroids, PYR) (Hongsibsong et al., 2019; Panuwet et al., 2009). Recently, 3-phenoxybenzoic acid metabolite of PYRs was found in serum samples from farmers and consumers (Thiphom et al., 2014). Furthermore, organochlorine pesticides, persistent compounds, were frequently detected in serum and breast milk (Stuetz et al., 2001); OPs; non-persistent compounds, were also detected in maternal serum and breast milk samples (Naksen et al., 2014). These biological metabolites of the pesticides detected are evidence of the exposure to OP and PYR pesticides. Furthermore, there has been a dramatic increase in the import of pesticides into Thailand during 2002–2016 (Department of Agriculture, Office of Agricultural Economics, Thailand). It is appeared that herbicides, insecticides, and fungicides are among the major pesticides used in Thailand and that the total quantity of pesticides imported into Thailand increased about fourfold, from 2002 to 2016.

Due to the limitation of available analytical resources, data on pesticide residues in vegetables and fruits in Thailand are scarce. To our knowledge, there are no reports from Thailand of multiple pesticide residues in vegetables and fruits. Here we report the series' survey of multiple pesticide residues in 16 different vegetables ($n = 412$) and 11 different fruits ($n = 301$) collected from rural and urban markets from four provinces engaged in intensive agriculture in upper northern Thailand during 2007–2013.

MATERIALS AND METHODS

Study sites

Vegetable and fruit samples were bought from rural and urban markets in four provinces engaged in intensive agriculture in upper northern Thailand, including Chiang Mai ($18^{\circ}50'14''N$ $98^{\circ}58'14''E$) (the site it was most conducted), Chiang Rai

(19°54'N 99°49'E), Lampang (18°18'N 99°30'E), and Phayao (19°9'55"N 99°54'13"E) provinces, as shown in Figure 1.




Figure 1. A map of four provinces including Chiang Mai, Chiang Rai, Lampang, and Phayao which were the study sites in this study in year 2010.

Collecting and processing vegetable and fruit samples

Sixteen different vegetables and 11 different fruits (Table 1) that are commonly consumed were collected from urban and rural markets during 2007–2013.

Table 1. Sixteen vegetables and eleven fruits sampled from various markets during 2007–2013.










	Common name (Thai name)	Scientific name
Vegetables		
1	 Baby corn (Khoa Pod On)	<i>Zea mays</i>
2	 Broccoli	<i>Brassica oleracea</i>
3	 Cabbage (Ka Lum)	<i>Brassica oleracea</i> L. var. <i>capitata</i> L.
4	 Cauliflower (Ka Lum Dok)	<i>Brassica oleracea</i> L. var. <i>botrytis</i> L
5	 Chili pepper (Prik Chi Fah)	<i>Capsicum annum</i>
6	 Chinese cabbage (Pak Kad Kow)	<i>Brassica pekinensis</i>
7	 Pakchoi (Pak Kwang Tung)	<i>Brassica chinensis</i> Just var <i>parachinensis</i> (Bailey) Tsen & Lee
8	 Chinese kale (Ka Nah)	<i>Brassica alboglabra</i>
9	 Cucumber (Tang Kwa)	<i>Cucumis sativus</i>

Table 1. Continued

	Common name (Thai name)	Scientific name
Vegetables		
10	 Ginger (Khing)	<i>Zingiber officinale</i>
11	 Bird's chili, Thai chili (Prik Ki Noo)	<i>Capsicum frutescens</i>
12	 Pepper (Prik Tai)	<i>Piper nigrum</i>
13	 Sugar pea, Sweet pea, Garden pea (Tua Lan Tao)	<i>Pisum sativum</i>
14	 Tomato (Ma Khua Thet)	<i>Lycopersicon esculentum</i>
15	 Water spinach, Water morning glory (Pak Bung)	<i>Ipomoea aquatica</i>
16	 Yard long bean (Tua Fak Yao)	<i>Vigna unguiculata</i>
Fruits		
17	 Apple	<i>Malus domestica</i>
18	 Asian pear (Sa Lhi)	<i>Pyrus pyrifolia</i>
19	 Dragon fruit (Keaw Mang Kon)	<i>Hylocereus undatus</i>
20	 Guava (Fa Rang)	<i>Psidium guajava</i>
21	 Grape (A Ngoon)	<i>Vitis vinifera</i>
22	 Lychee (Lin Jee)	<i>Litchi chinensis</i>
23	 Mango (Ma Muang)	<i>Mangifera indica</i>
24	 Rose apple (Chom Phoo)	<i>Syzygium samarangense</i>
25	 Strawberry	<i>Fragaria ananassa</i>
26	 Tangerine (Som Kheaw Wan)	<i>Citrus Reticulata</i>
27	 Watermelon (Tang Mo)	<i>Citrullus lanatus</i>

Sample collection

During 2007–2013, Chiang Mai urban (local and supermarkets) and rural areas were major sites of vegetable and fruit sample collection for pesticide residue analysis in the present study. However, since Chiang Mai province was a major source of tangerine cultivation in the country, contamination of tangerine with pesticide residues

was a public health concern (Prapamontol et al, 2010). Therefore, in 2009, 100 fresh tangerine samples were collected from rural markets in Chiang Mai for multiple pesticide residue analysis. Besides the studies on Chiang Mai province, in 2010, the study sites were extended to three more provinces of upper northern Thailand practicing intensive agriculture. Therefore, in 2010, vegetable and fruit samples were collected from rural markets in Chiang Mai, Chiang Rai, Lampang, and Phayao provinces. Furthermore, in 2013, vegetables and some medicinal plants, that is, ginger and pepper, were analyzed for two groups of fungicides, namely carbendazim and dithiocarbamates. The details of the sample collection in each of the years, sites, pesticide residues, and number of samples are shown in Table 2.

The methods adopted for sampling the vegetables and the fruits were based on Codex recommended methods of sampling for the determination of pesticide residues for compliance with MRLs (CODEX, 1999).

For sampling vegetables and fruits from different markets, each type of vegetable and fruit was purchased from four or more retail shops (4 units) in each market. Approximately, 1 kg of each vegetable or fruit type was purchased from one retail shop. Each unit was divided into four portions. One portion was put in sterile polythene bags with label and stored on ice in a Styrofoam box during transportation to the toxicology laboratory of the Environment and Health Research Unit, Research Institute for Health Sciences, Chiang Mai University. All the labeled samples were stored at -20°C until analysis. The diagram of the sampling protocol is presented in Figure 2. All samples were not washed and cleaned according to CODEX (1999) Edible parts of all the samples were chopped into small pieces and homogenized before the extraction procedure.

Table 2. Details of vegetable and fruit sample collection during 2007–2013.

Year	2007	2008	2009	2010	2013
Sampling site					
-Areas	Urban	Urban	Rural	Rural	Urban
-Provinces (n)	Chiang Mai (150)	Chiang Mai (150)	Chiang Mai (156)	Chiang Mai (53) Chiang Rai (39) Lampang (41) Phayao (38)	Chiang Mai (86)
Types (n)	-Vegetables (100) -Fruits (50)	-Vegetables (100) -Fruits (50)	-Vegetables (35) -Fruits (21) -Tangerines (100)	-Vegetables (91) -Fruits (80)	-Vegetables (86)
Analyzed pesticides	OP pesticides	OP pesticides	- OP pesticides -Carbamates -Synthetic pyrethroids -Carbendazim -Imidacloprid -Abamectins	OP pesticides	-Carbendazim -Dithiocarbamates




Figure 2. A sampling protocol for the different types of vegetables and fruits from one market.

Chemicals and standards

All the organic solvents were of analytical grade. Acetone, acetonitrile, ethyl acetate, and toluene were purchased from J.T. Baker (Radner, Pennsylvania, USA). Sodium chloride, sodium sulfate, potassium dihydrogen phosphate, and dipotassium hydrogen phosphate were purchased from Merck (Darmstadt, Germany). The graphite carbon black (GCB) was purchased from Agilent Technologies (USA). The standard pesticides: 20 organophosphate (OP) pesticides, 10 carbamates, and 5 synthetic pyrethroids, and bifenthrin, carbendazim, imidacloprid, abamectin, and imidacloprid were purchased from Dr. Ehrenstorfer GmbH (Augsburg, Germany). Triphenylphosphate was purchased from Fluka (Buchs, Germany).

Methods

The pesticides selected for the determination of the presence of pesticide residues in the samples were based on the data on pesticides imported into Thailand (OAE, 2013). An aliquot of sample was analyzed using internal standard and matrix-matched standard calibration curve. Quality assurance and control was performed using spiked matrix-matched aliquot (in duplicate) as a sample. Pesticide analysis methods for OPs (Polyiem et al., 2018), carbamates (Tuan et al., 2009), synthetic PYRs (Pakvilai et al., 2015), abamectins (Diserens and Henzelin, 1999), carbendazim, dithiocarbamates, and imidacloprid were performed according to methods previously described. The methods employed in this study have limit of detection (LOD) and limit of quantitation (LOQ) below the MRLs set by *Codex Alimentarius*. In case Codex MRLs were not available, MRLs from Thailand (Thai Agricultural Commodity and Food Standard TACFS 9002-2008), Taiwan (<https://law.moj.gov.tw/ENG/>), Korea (<http://www.foodsafetykorea.go.kr/residue/prd/>), Japan (<http://db.ffcr.or.jp/front/>), and the EU (Draft EU MRLs in view of the first establishment of Annex II to Regulation (EC) N.396/2005) were adopted. Each batch of sample analysis had at least an aliquot of spiked pooled matrix added as the internal quality control sample.

Organophosphate pesticides. OP pesticides have been widely used in vegetable and fruit cultivation. The method used for detecting 20 OP pesticide residues was as described previously (Polyiem et al., 2018). Twenty OP pesticides analyzed were azinphos-ethyl, azinphos-methyl, chlorpyrifos, diazinon, dicrotophos, dimethoate, EPN,

ethion, fenitrothion, malathion, methamidophos, methidathion, mevinphos, monocrotophos, parathion-methyl, pirimiphos-ethyl, pirimiphos-methyl, profenofos, prothiophos, and triazophos. Briefly, 5 g of well-homogenized vegetable and fruit samples was extracted with 5 mL acetonitrile and cleaned using 500 mg/5 mL graphite carbon black. The ethyl acetate extract (1 μ L) was analyzed by gas chromatography–flame photometric detection (Hewlett Packard 6890 Series, Agilent Technologies, CA, USA).

Cabamate pesticides. The method used for detecting the 10 carbamate pesticides was adopted from the Taiwan Agricultural Chemicals and Toxic Substances Research Institute, Taichung, Taiwan (Tuan et al., 2009). Ten carbamate pesticides assayed were 3-hydroxy carbofuran, aldicarb, aldicarb sulfone, aldicarb sulfoxide, Baygon, carbaryl, carbofuran, methiocarb, methomyl, and oxamyl. Briefly, 10 g of the homogenized sample was extracted with acetonitrile and then cleaned. The clean acetone extract was subjected to post-column derivatization with o-phthalaldehyde (OPA) and analyzed using high performance liquid chromatography–fluorescence detection (Shimadzu SCL-10A VP, Shimadzu, Japan).

Synthetic pyrethroids pesticides. Since PYR pesticides have enantiometric isomers (Corcellas et al., 2015), the present study used the method reported by Pakvilai et al. (Pakvilai et al., 2015), and the summation of the area peaks from each PYR pesticide was employed for the calculation of the concentration. Six synthetic PYR pesticides including cyfluthrin, cypermethrin, deltamethrin, fenvalerate, lambda cyhalothrin, and permethrin were analyzed. In brief, 5 g of the homogenized sample was extracted with 10 mL of dichloromethane. The filtered extracts were cleaned using 250 mg graphitized carbon cartridge solid phase extraction. The clean extract was analyzed using gas chromatography–electron capture detection (Hewlett Packard 6890 Series, Agilent Technologies, CA, USA).

Carbendazim. Analysis of carbendazim in vegetable samples was carried out by employing the method reported by Phansawan et al. (2015). In brief, 5 g of the homogenized vegetable sample was extracted with ethyl acetate in acidic condition. The extract was cleaned by filtering through a SAX/PSA dual-layer cartridge. The clean extract was dissolved in HPLC mobile phase (methanol:water, 25:75, v/v) and analyzed using HPLC-variable UV detection (Agilent 1100, Agilent Technologies, CA, USA).

Dithiocarbamates. The method used for analysis of DTCs in vegetables such as cucumber, ginger, and pepper was a method reported by Phansawan et al. (2014). Briefly, 5 g of homogenized fresh vegetable samples was transferred into the extraction tube, followed by the addition of 4.0 mL of freshly prepared 0.5% SnCl₂ made in 5 M hydrochloric acid. Then, into each tube was added 1.0 mL of isooctane. The samples were heated in an ultrasonic water bath at 80°C for 40 min with continuous stirring and then allowed to cool down to room temperature (25°C). During cooling, the tubes were periodically shaken by hand to assist with the CS₂ sipping into the isooctane layer, and 1 μ L of the isooctane layer was injected into a gas chromatography–flame photometric detector (Hewlett Packard 6890 Series, Agilent Technologies, CA, USA).

Imidacloprid. The analysis of imidacloprid was carried out by using a method modified from the method reported previously (Ferrer and Thurman, 2007). In brief, 10 g of the homogenized sample was extracted by using acetonitrile (1% acetic acid), and the extract was analyzed using HPLC-UV detection (Agilent 1100, Agilent Technologies, CA, USA).

Abamectins. Analysis of abamectins in fruits was performed using the method reported by Discerns and Henzelin with slight modifications (Diserens and Henzelin, 1999). In brief, 10 g of the homogenized sample was extracted using methanol and cleaned by solid phase extraction (Vertipak C18 SPE, Vertical, Bangkok, Thailand). The extract was derivatized with reagent 1 (acetonitrile and trifluoroacetic acid, 2:1) and reagent 2 (acetonitrile and 1-methylimidazole, 1:1), and analyzed by HPLC-Fluorescence detection (Agilent 1100, Agilent Technologies, CA, USA).

RESULTS

This study reported pesticide residues in vegetables and fruits from four provinces in northern Thailand during 2007–2013. A total of 713 vegetable and fruit samples were analyzed for 43 multi-classes of pesticides including insecticides (20 OPs, 10 carbamates, 6 PYRs, 2 abamectins, and imidacloprid), and fungicides (carbendazim and dithiocarbamate). However, organochlorine pesticides were not included in this study because they have been banned in agriculture by the Thai government since 1983 (IPM, 2015).

Multi-pesticide residues found in vegetable and fruit samples

Table 3 summarizes the incidence of pesticide residues found in vegetables and fruits from rural and urban markets. Individual pesticide residues detected and their concentrations (mg/kg) in vegetables and fruits can be found in Supplement A.

Table 3. Summary of pesticide residues found in vegetables and fruits from rural and urban markets in upper northern Thailand during 2007-2013.

Types	Number of pesticide residues detected	Urban markets	Rural markets
Vegetables	Single & multiple	42.6%	63.5%
	Multiple	16.8%	29.4%
		<i>N</i> = 286	<i>N</i> = 126
Fruits	Single & multiple	74%	83.3%
	Multiple	38%	67.2%
		<i>N</i> = 100	<i>N</i> = 201

It was appeared that vegetable and fruit samples bought from rural markets had pesticide residues with higher frequency of detection, in both 'single & multiple' and 'multiple', than vegetable and fruit samples from urban markets.

The fruit samples were found to have more residue-detected samples than the vegetable samples. The vegetable and the fruit samples with the highest percentages of contamination were, respectively, Chinese kale (74%) and tangerine (98%). There were 21 different pesticides detected in the tangerine samples. Baby corn was the only type of sample in this study which was not found to have pesticide contamination.

Frequently detected pesticides

All the pesticides and their percentages of detection are shown in Supplement B. Among the 20 OPs analyzed in the vegetable samples, chlorpyrifos was the most-detected OP. Twenty percent of the samples from urban markets and 18.9% of the samples from rural markets had chlorpyrifos levels that exceeded the MRL. Among 10 carbamates, aldicarb and carbaryl were found in the fruits (tangerines), and exceeded the MRLs by 1% and 2%, respectively. Synthetic PYRs were the most frequently detected residue in the both vegetables and fruits samples from rural markets and all six compounds exceeded the MRLs by values between 2.9% and 65.5%. Abamectin was found only in the tangerine samples, and only abamectin 1a compound was found to exceed the MRL in 13% of the analyzed samples. Cabendazim and imidacloprid were also found in the tangerine samples, but their contents did not exceed the MRL. However, in the vegetable samples from urban markets, the carbendazim residues detected were found to exceed the MRL by 15.6%. Dithiocarbamate fungicide was also detected, and found to exceed the MRL by 5.6%. In summary, among the 43 compounds of pesticides investigated in the present report, 27 (62.8%) were detected in the vegetable and fruit samples. According to the present report, the top 10 pesticide

residues that exceeded the MRL show as following: (1) lambda-cyhalothrin (65.5%), (2) deltamethrin (56.7%), (3) cyfluthrin (34.5%), (4) fenvalerate (29.8%), (5) cypermethrin (21.6%), (6) chlorpyrifos (19.6%), (7) profenophos (5.4%), (8) methamidophos (5.2%), (9) ethion (4.8%), and (10) permethrin (2.9%).

Comparison of pesticide residues found in four provinces

In 2010, 171 vegetable and fruit samples from 4 provinces including Chiang Mai, Chiang Rai, Lampang, and Phayao were analyzed for multi-pesticide residues. The number of OP pesticide-contaminated samples sourced from Chiang Mai was the highest (Figure 3A). At the same time, the numbers of pyrethroid pesticide-contaminated samples obtained from the four provinces were not markedly different (Figure 3B).




Figure 3. Percentages of OP pesticide (A) and pyrethroid (B) residue-detected samples (both below and above MRLs) in Chiang Mai Province during 2007–2010, and the percentages in the four provinces in 2010.

DISCUSSION

The present report shows that multiple pesticide residues were detected in vegetable and fruit samples with high frequency. The frequencies of pesticide residues detected were higher in samples purchased from rural markets than in samples from urban markets. Similar reports showed higher pesticide residue concentrations in vegetables collected from the farms than bought from the markets and supermarkets (Sapbamrer and Hongsibsong, 2014; Kumari and John, 2019). This phenomenon might be explained from recent or just after cultivated produce as most of common consume

vegetable and fruits growing in the rural areas. Furthermore, lower residue detection in vegetables sold urban markets and supermarkets may be explained by pre-processing step which include washing and removal of unwanted parts. Overall, pesticide residues were more frequently detected in fruits than in vegetables (Table 3). In addition, among samples of vegetables ($n = 412$) and fruits ($n = 301$), 45% of vegetable and 55% of fruit samples found pesticide residues exceeded their MRLs (Table 4). Among 43 pesticides analyzed, residues of pyrethroid pesticides detected in samples from four provinces in 2010 almost all exceeded the MRLs (Figure 3B) compared with residues of OP pesticides (Figure 3A). Chlorpyrifos, an OP pesticide, was the most detected OP in the present report and among the top 10 in the list of active ingredients imported into Thailand during 2007-2014 (OAE, 2013). Chlorpyrifos is a common OP pesticide detected in vegetables and fruits in Thailand. It has been reported to be found in vegetables from markets around Kwan Phayao Lake, Phayao Province, upper northern Thailand (Sapbamrer and Hongsibsong, 2014), Chinese kale (purchased from markets in Nakhon Pathom Province, central Thailand (Wanwimolruk et al., 2015), and mangosteen bought from markets in Bangkok, central and eastern provinces of Thailand (Phopin et al., 2017) were also found to contain chlorpyrifos residues.

MRL is the maximum concentration of a pesticide residue (expressed as mg/kg), recommended by the Codex Alimentarius Commission to be legally permitted in or in food commodities and animal feeds. MRLs are based on good agricultural practices (GAP) data and foods derived from commodities that comply with the respective MRLs and are intended to be toxicologically acceptable (<http://www.fao.org/waicent/faostat/Pest-Residue/pest-e.htm>). In order to protect consumers' health, residues of pesticides in any foods must not above or exceed their MRLs.

This report is derived from survey series of pesticide residues for 5 years in 2007, 2008, 2009, 2010 and 2013. To our best knowledge, this is the longest survey duration ever reported from Thailand. Furthermore, this study covered several different pesticide groups used in Thailand namely organophosphates, carbamates, synthetic pyrethroids, carbendazim, dithiocarbamates, imidaclopid, abamectins which constitute 43 pesticides in total. This report also covered vegetables and fruits that common consume in northern as well as other parts of Thailand. Zaidon et al. (2016) reported a review on pesticides occurrence in fruits and vegetables in Malaysia from 2007 to 2014 covered 10 pesticides and 30% of pesticide residues detected exceeded their MRLs.) Poulsen et al. (2017) also reported the results from the Danish monitoring programme for pesticide residues from 2004-2011.

This survey was conducted in the academic institution with some limitation of funding for some time over a long period of time (2007-2013). In order to protect consumers' health, national institution (s) would be able to secure funding over years' plan such as Thai Ministry of Public Health by regular monitoring of pesticide residues in foods and Ministry of Agriculture and Cooperatives by enforcement monitoring of good agricultural practices (GAP).

CONCLUSION

A total of 713 samples of vegetables and fruits collected from urban and rural markets in northern Thailand during 2007-2013 were analyzed for 43 pesticide residues. This study found that 67% of the vegetable and fruit samples had pesticide residues and 43% had residues that exceeded the MRLs set by Codex and others, namely the EU, Japan, Taiwan, Korea, and Thailand.

The fruit samples were found to have percentage of pesticide detection higher than the vegetable samples. Of the 301 fruit samples, 81% contained pesticide residues and 55% exceeded the MRL while in the 412 vegetable samples, 57% had pesticide residues and 45% exceeded the MRL. The samples collected from rural markets had percentage detection of pesticide residues higher than the samples collected from urban markets. The top 10 of the pesticide residues detected in the present report include 6 synthetic pyrethroids and 4 organophosphate pesticides. Lambda-cyhalothrin was the most frequently detected synthetic pyrethroid and chlorpyrifos was the most frequently detected organophosphate pesticide.

Table 4. Summary of pesticide residue detection in vegetable and fruit samples from urban and rural markets in upper northern Thailand during 2007-2013.

Site	Total samples	Detection in sample (% of total samples)	Exceeding MRL (% of total samples)	Type of pesticide	No. of analyzed samples	Detection in sample (% of analyzed samples)	Exceeding MRL (% of analyzed samples)	Detected pesticides
All	Total (N = 713)	480 (67%)	308 (43%)					
	Vegetables (n = 412)	235 (57%)	185 (45%)					
	Fruits (n = 301)	245 (81%)	165 (55%)					
Urban (n = 386)	Total (n = 386)	199 (52%)	96 (25%)					
	Vegetables (n = 286)	125 (44%)	58 (20%)	OP pesticides	200	94 (47%)	50 (25%)	azinophos-methyl, chlorpyrifos, diazinon, dicrotophos, dimethoate, EPN, ethion, malathion, methamidophos, mevinphos, monocrotophos, parathion-methyl, pirimiphos-ethyl, pirimiphos-methyl, profenophos, triazophos (n = 16)
				carbamates	Not analyzed			-
				pyrethroids	Not analyzed			-
				others	86	31 (36%)	8 (9%)	carbendazim, mancozeb (n=2)
	Fruits (n = 100)	74 (74%)	38 (38%)	OP pesticides	100	74 (74%)	38 (38%)	chlorpyrifos, diazinon, dimethoate, ethion, fenitrothion, malathion, methamidophos, methidathion, monocrotophos, parathion-methyl, pirimiphos-ethyl, pirimiphos-methyl, profenophos, prothiophos, triazophos (n = 15)
				carbamates	Not analyzed			-
				pyrethroids	Not analyzed			-
				others	Not analyzed			-

Table 4. Continued.

Site	Total samples	Detection in sample (% of total samples)	Exceeding MRL (% of total samples)	Type of pesticide	No. of analyzed samples	Detection in sample (% of analyzed samples)	Exceeding MRL (% of analyzed samples)	Detected pesticides
Rural (n=327)	Total (n = 327)	281 (86%)	212 (65%)					
	Vegetables (n = 126)	110 (87%)	85 (67%)	OP pesticides	126	110 (87%)	29 (26%)	chlorpyrifos, diazinon, dicotophos, dimethoate, parathion-methyl, ethion, malathion, methamidophos, monocrotophos, parathion-methyl, profenophos, triazophos (n = 12)
				carbamates	Not analyzed			-
				pyrethroids	91	85 (93%)	85 (93%)	cyfluthrin, cypermethrin, deltamethrin, fenvalerate, lamda-cyhalothrin, permethrin (n = 6)
			others	Not analyzed			-	
	Fruits (n = 201)	171 (85%)	127(63%)	OP pesticides	201	171 (85%)	51 (25%)	chlorpyrifos, diazinon, dicotophos, dimethoate, ethion, malathion, methamidophos, monocrotophos, profenophos, prothiophos, triazophos (n = 11)
				carbamates	100	1 (1%)	1 (1%)	abamectin 1a, aldicarb, carbaryl,
				pyrethroids	180	159 (88%)	127 (70%)	cyfluthrin, cypermethrin, deltamethrin, fenvalerate, lamda-cyhalothrin (n = 6)
				others	100	87 (87%)	4 (4%)	carbendazim, imidacloprid

This study result may add to the existing evidence of pesticide residues which could be able to estimate adverse effect on public health for policy maker. Finally, multi-residue methods are proposed as a regular monitoring system to enhance food safety policy both internal consumption and exportation.

REFERENCES

- Arias, L.A., Bojacá, C.R., Ahumada, D.A., and Schrevens, E. 2014. Monitoring of pesticide residues in tomato marketed in Bogota, Colombia. *Food Control*. 35:213-217.
- Boobis, A.R., Ossendorp, B.C., Banasiak, U., Hamey, P.Y., Sebestyen, I., and Moretto, A. 2008. Cumulative risk assessment of pesticide residues in food. *Toxicology Letters*. 180: 137-150.
- Chalermphol, J., Bastakoti, G.B., and Bastakoti, R.C. 2014. Adoption of improved varieties of vegetable crops with pesticide use in Chiang Mai Province, Northern Thailand. *Procedia Environmental Sciences*. 20: 418-424.
- CODEX. 1999. Recommended methods of sampling for the determination of pesticide residues for compliance with MRLs. CAC/GL33-1999.
- Corcellas, C., Eljarrat, E., and Barcelo, D. 2015. Enantiomeric-selective determination of pyrethroids: application to human samples. *Analytical and Bioanalytical Chemistry*. 407: 779-786.
- Diserens, H., and Henzelin, M. 1999. Determination of abamectin residues in fruits and vegetables by high-performance liquid chromatography. *Journal of Chromatography A*. 833: 13-18.
- Ferrer, I., and Thurman, E.M. 2007. Multi-residue method for the analysis of 101 pesticides and their degradates in food and water samples by liquid chromatography/time-of-flight mass spectrometry. *Journal of Chromatography A*. 1175: 24-37.
- Hjorth, K., Johansen, K., Holen, B., Andersson, A., Christensen, H.B., Siivinen, K., and Toome, M. 2011. Pesticide residues in fruits and vegetables from South America – A Nordic project. *Food Control*. 22: 1701-1706.
- Hongsibsong, S., Prapamontol, T., Dong, J.X., Bever, C.S., Xu, Z.L., Gee, S.J., and Hammock, B.D. 2019. Exposure of consumers and farmers to organophosphate and synthetic pyrethroid insecticides in Northern Thailand. *Journal fur Verbraucherschutz und Lebensmittelsicherheit*. 14: 17-23.
- Jardim, A.N.O., and Caldas, E.D. 2012. Brazilian monitoring programs for pesticide residues in food – Results from 2001 to 2010. *Food Control*. 25: 607-616.
- Kumari, D., and John, S. 2019. Health risk assessment of pesticide residues in fruits and vegetables from farms and markets of Western Indian Himalayan region. *Chemosphere*. 224: 162-167.
- Naksen, W., Prapamontol, T., Mangklabruks, A., Chantara, S., and Thavornytikarn, P. 2014. Health risk from pesticide exposure among infants of breastfeeding farm workers in an agricultural area of Chiang Mai Province, Thailand. *Advances in Environmental Biology*. 8: 213-218.
- National Statistical Office, M.o.I.a.C.T. 2003. Thailand Agricultural Census.
- Nougadère, A., Sirot, V., Kadar, A., Fastier, A., Truchot, E., Vergnet, C., Hommet, F., Baylé, J., Gros, P., Leblance, J.-C. 2012. Total diet study on pesticide residues in France: Levels in food as consumed and chronic dietary risk to consumers. *Environment International*. 45: 135-150.
- Pakvilai, N., Prapamontol, T., Thavornytikarn, P., Mangklabruks, A., Chantara, S., Hongsibsong, S., and Santasup, C. 2015. A simple and sensitive GC-ECD method for detecting synthetic pyrethroid insecticide residues in vegetable and fruit samples. *Chiang Mai Journal of Science*. 42: 197-208.
- Panseri, S., Catalano, A. Giorgi, A., Arioli, F., Procopio, A., Britti, D., and Chiesa, L.M. 2014. Occurrence of pesticide residues in Italian honey from different areas in relation to its potential contamination sources. *Food Control*. 38: 150-156.

- Panuwet, P., Prapamontol, T., Chantara, S., and Barr, D.B. 2009 Urinary pesticide metabolites in school students from northern Thailand. *International Journal of Hygiene and Environmental Health*. 212: 288-297.
- Panuwet, P., Prapamontol, T., Chantara, S., Thavornyuthikarn, P., Montesano, M.A., Whitehead, R.D. Jr., and Barr, D.B. 2008. Concentrations of urinary pesticide metabolites in small-scale farmers in Chiang Mai Province, Thailand. *The Science of The Total Environment*. 407: 655-668.
- Phansawan, B., Prapamontol, T., Chantara, S., Mangklabruks, A., Santasup, C., and Thavornyutikarn, P. 2014. Determination of dithiocarbamate fungicide residue in cucumber, ginger, and pepper and health risk assessment. *Advances in Environmental Biology*. 8: 45-50.
- Phansawan, B., Prapamontol, T., Thavornyutikarn, P., Chantara, S., Mangklabruks, A., and Santasup, C. 2015. A sensitive method for determination of carbendazim residue in vegetable samples using HPLC-UV and its application in health risk assessment. *Chiang Mai Journal of Science*. 42: 681-690.
- Phopin, K., Wanwimolruk, S., and Prachayasittikul, V. 2017. Food safety in Thailand. 3: Pesticide residues detected in mangosteen (*Garcinia mangostana* L.), queen of fruits. *Journal of the Science of Food and Agriculture*. 97: 832-840.
- Polyiem, W., Naksen, W., and Prapamontol, T. 2018. Gas chromatographic-flame photometric detection of organophosphate pesticide residues and its application in real vegetable and fruit samples from Chiang Mai city, Thailand. *Chiang Mai Journal of Science*. 45: 1933-1943.
- Pose-Juan, E., Sánchez-Martín, M.J., Andrades, M.S., Rodríguez-Cruz, M.S., and Herrero-Hernández, E. 2015. Pesticide residues in vineyard soils from Spain: Spatial and temporal distributions. *Science of The Total Environment*. 514: 351-358.
- Posri, W., Shankar, B., and Chadbunchachai, S. 2006. Consumer attitudes towards and willingness to pay for pesticide residue limit compliant 'safe' vegetables in Northeast Thailand. *Journal of International Food & Agribusiness Marketing*. 19: 81-101.
- Poulsen, M.E., Andersen, J.H., Petersen, A., and Jensen, B.H. 2017. Results from the Danish monitoring programme for pesticide residues from the period 2004–2011. *Food Control*. 74: 25-33.
- Prapamontol, T., Hongsibsong, S., Pakvilai, N., Polyiem, W., Kawichai, S., and Santasup, C. 2010. Pesticide residues in tangerines (*Citrus reticulata* Blanco) cultivated in different types from Chiang Mai Province, northern Thailand. *Toxicology Letters*. 196 Supple: S334-S335.
- Riwthong, S., Schreinemachers, P., Grovermann, C., and Berger, T. 2015. Land use intensification, commercialization and changes in pest management of smallholder upland agriculture in Thailand. *Environmental Science & Policy*. 45: 11-19.
- Sangchan, W., Hugenschmidt, C., Ingwersen, J., Schwadorf, K., Thavornyutikarn, P., Pansombat, K., and Streck, T. 2012. Short-term dynamics of pesticide concentrations and loads in a river of an agricultural watershed in the outer tropics. *Agriculture, Ecosystems & Environment*. 158: 1-14.
- Sapbamrer, R., and Hongsibsong, S. 2014. Organophosphorus pesticide residues in vegetables from farms, markets, and a supermarket around Kwan Phayao Lake of Northern Thailand. *Archives of Environmental Contamination and Toxicology*. 67: 60-67.
- Schreinemachers, P., Sringarm, S., and Sirijinda, A. 2011. The role of synthetic pesticides in the intensification of highland agriculture in Thailand. *Crop Protection*. 30: 1430-1437.

- Schreinemachers, P., and Tipraqsa, P. 2012. Agricultural pesticides and land use intensification in high, middle and low income countries. *Food Policy*. 37: 616-626.
- Skretteberg, L.G., Lyrån, B., Holen, B., Jansson, A., Fohgelberg, P., Siivinen, K., Andersen, J.H., and Jensen, B.H. 2015. Pesticide residues in food of plant origin from Southeast Asia – A Nordic project. *Food Control*. 51: 225-235.
- Stuetz, W., Prapamontol, T., Erhardt, J.G., and Classen, H.G. 2001. Organochlorine pesticide residues in human milk of a Hmong hill tribe living in Northern Thailand. *Science of the Total Environment*. 273: 53-60.
- Szpyrka, E., Kurdziel, A., Matyaszek, A., Podbielska, M., Rupa, J., and Słowik-Borowiec, M. 2015. Evaluation of pesticide residues in fruits and vegetables from the region of south-eastern Poland. *Food Control*. 48: 137-142.
- Thiphom, S., Prapamontol, T., Chantara, S., Mangklabruks, A., Suphavitai, C., Ahn, K.C., Gee, S.J., and Hammock, B.D. 2014. Determination of the pyrethroid insecticide metabolite 3-PBA in plasma and urine samples from farmer and consumer groups in northern Thailand. *Journal of Environmental Science and Health. Part. B, Pesticides, Food Contaminants, and Agricultural Wastes*. 49: 15-22.
- Tuan, S.J., Tsai, H.M., Hsu, S.M., and Li, H.P. 2009. Multiresidue analysis of 176 pesticides and metabolites in pre-harvested fruits and vegetables for ensuring food safety by gas chromatography and high performance liquid chromatography. *Journal of Food and Drug Analysis*. 17: 163-177+230.
- Wanwimolruk, C., Phopin, K., and Wanwimolruk, S. 2019. Food safety in Thailand 6: How to eat guava fruits safely? Effects of washing and peeling on removing pesticide residues in guava fruits. *Journal of Food Safety*. 39.
- Wanwimolruk, S., Kanchanamayoon, O., Phopin, K., and Prachayasittikul, V. 2015. Food safety in Thailand 2: Pesticide residues found in Chinese kale (*Brassica oleracea*), a commonly consumed vegetable in Asian countries. *Science of the Total Environment*. 532: 447-455.
- Wongta, A., Sawarng, N., Tongchai, P., Sutan, K., Kerdnoi, T., Prapamontol, T., and Hongsibsong, S. 2018. The pesticide exposure of people living in agricultural community, Northern Thailand. *Journal of Toxicology*. 2018(6): 1-7.
- Zaidon, S.Z., Hamsan, H., and Bin, H.Y. 2016. A review on pesticides occurrence in fruits and vegetables in Malaysia and their potential health risk among adults. *Indian Journal of Environmental Protection*. 36: 826-832.

OPEN access freely available online

Chiang Mai University Journal of Natural Sciences [ISSN 16851994]

Chiang Mai University, Thailand

www.cmuj.cmu.ac.th

Supplement A. Pesticide residues detected in vegetable and fruit samples in upper northern Thailand during 2007–2013.

Samples	Urban					Rural				
	N	No. of contaminated samples	No. of samples detected multi-residues	Detected pesticides; range or mean ^b (mg/kg)	No. exceeding MRL	N	No. of contaminated samples	No. of samples detected multi-residues	Detected OP pesticides; range or mean ^b (mg/kg)	No. exceeding MRL
Baby Corn	20	0	0	none	0	-	-	-	-	-
Broccoli	20	12	4	chlorpyrifos (0.005–0.030) dicotophos (0.014) diazinon (0.014) ethion (0.002) mevinphos (0.030–0.011) pirimiphos-ethyl (0.008–0.018) pirimiphos-methyl (0.005)	2 0 0 0 0 0 0	-	-	-	-	
Cabbage	20	5	3	diazinon (0.009–0.402) azinophos-methyl (0.088)	0 0	24	11	3	chlorpyrifos (0.002–0.020) dicotophos (0.010) ethion (0.001) methamidophos (0.015–0.017) monocrotophos (0.044) cyfluthrin (0.047–0.656) cypermethrin (0.028–0.321) deltamethrin (0.079–4.854) fenvalerate (0.131–2.374) lamda-cyhalothrin (0.063–0.762) permethrin (0.002–0.063)	1 0 0 2 0 9 2 13 12 14 0
Cauliflower	20	10	8	chlorpyrifos (0.002–0.232) dimethoate (0.004) ethion (0.001) methamidophos (0.005–0.639) pirimiphos-methyl (0.013)	7 0 0 4 0	9	3	0	chlorpyrifos (0.001–0.016) dicotophos (0.027)	1 0
	4	3	NA	carbendazim (0.013–0.020)	1				cyfluthrin (0.028–0.276) cypermethrin (0.031–0.402) deltamethrin (0.436–2.002) fenvalerate (0.129–0.867) lamda-cyhalothrin (0.185–0.757) permethrin (0.074–0.261)	6 1 6 3 8
Chili pepper	4	3	NA	carbendazim (0.038–0.040)	0	-	-	-	-	-

Supplement A. Continued

Samples	Urban					Rural				
	N	No. of contaminated samples	No. of samples detected multi-residues	Detected pesticides; range or mean ^b (mg/kg)	No. exceeding MRL	N	No. of contaminated samples	No. of samples detected multi-residues	Detected OP pesticides; range or mean ^b (mg/kg)	No. exceeding MRL
Chinese cabbage	20	7	2	chlorpyrifos (0.009–0.043)	5	15	12	9	chlorpyrifos (0.001–1.596)	3
				diazinon (0.001)	0				dicrotophos (0.001–0.139)	0
				pirimiphos-methyl (0.011–0.034)	0				diazinon (0.001–0.002)	0
				profenophos (0.025)	0				dimethoate (0.002–0.159)	0
									parathion-methyl (0.002)	0
									profenophos (0.001–0.055)	1
									triazophos (0.001)	0
									cyfluthrin (0.077–0.544)	5
									cypermethrin (0.051–7.728)	4
									deltamethrin (0.298–1.052)	6
Pakchoi	-	-	-		-	14	13	10	chlorpyrifos (0.001–1.596)	4
									dicrotophos (0.001–0.775)	1
									methamidophos (0.001–0.012)	1
									monocrotophos (0.001–0.313)	2
									parathion-methyl (0.020)	0
									profenophos (0.002–0.256)	1
									triazophos (0.006)	0
									cyfluthrin (0.060–0.525)	4
									cypermethrin (0.041–8.389)	3
									deltamethrin (0.404–4.753)	9
Chinese kale	20	13	11	chlorpyrifos (0.007–1.394)	1	18	15	9	chlorpyrifos (0.002–0.416)	3
				diazinon (0.144)	9				dicrotophos (0.020–0.973)	2
				dicrotophos (0.003–0.188)	1				ethion (0.001)	0
				ethion (0.001–0.020)	1				malathion (0.002)	0
				monocrotophos (0.003)	0				methamidophos (0.002)	0
				profenophos (0.013–0.130)	0				profenophos (0.160–0.251)	2

Supplement A. Continued.

Samples	Urban					Rural				
	N	No. of contaminated samples	No. of samples detected multi-residues	Detected pesticides; range or mean ^b (mg/kg)	No. exceeding MRL	N	No. of contaminated samples	No. of samples detected multi-residues	Detected OP pesticides; range or mean ^b (mg/kg)	No. exceeding MRL
Chinese kale	4	2	NA	carbendazim (0.034–0.119)	0				cyfluthrin (0.014–0.470)	6
									cypermethrin (0.044–0.594)	4
									deltamethrin (0.319–9.972)	7
									fenvalerate (0.028–2.583)	6
									lamda-cyhalothrin (0.038–0.472)	5
									permethrin (0.015–0.210)	0
Cucumber	20	13	6	chlorpyrifos (0.002–0.106)	6	19	12	2	chlorpyrifos (0.003–0.116)	4
				dimethoate (0.010)	0				dicotophos (0.010–0.034)	0
				malathion (0.006)	0				ethion (0.001)	0
				pirimiphos-ethyl (0.006–0.022)	0					
				pirimiphos-methyl (0.006–0.007)	0					
				profenophos (0.093)	1					
				carbendazim (0.013–0.020)	0				cyfluthrin (0.096–0.619)	6
				mancozeb (0.12–0.27)	0				cypermethrin (0.078–0.228)	1
									deltamethrin (0.434–0.681)	4
									fenvalerate (0.092–0.553)	2
		lamda-cyhalothrin (0.211–0.976)	4							
		permethrin (0.021–0.130)	0							
Ginger	24	2	NA	carbendazim (0.009–0.011)	0	-	-	-	-	-
Bird's chili	4	3	NA	carbendazim (0.011–0.081)	1	-	-	-	-	-
Pepper	14	7	NA	mancozeb (0.14–30.6)	3	-	-	-	-	-
Sweet pea	20	10	3	chlorpyrifos (0.005–0.649)	4	3	2	0	ethion (0.001)	0
				dimethoate (0.002–0.024)	0					
				methamidophos (0.015–0.032)	3					
				pirimiphos-methyl (0.005–0.006)	0					
Tomato	4	3	NA	carbendazim (0.014–0.064)	1					

Supplement A. Continued

Samples	Urban					Rural						
	N	No. of contaminated samples	No. of samples detected multi-residues	Detected pesticides; range or mean ^b (mg/kg)	No. exceeding MRL	N	No. of contaminated samples	No. of samples detected multi-residues	Detected OP pesticides; range or mean ^b (mg/kg)	No. exceeding MRL		
Water morning glory	20	9	6	chlorpyrifos (0.002–0.003)	0	8	4	0	chlorpyrifos (0.002–0.103)	2		
				dicrotophos (0.004–0.403)	2				dicrotophos (0.072)	0		
				ethion (0.001–0.002)	0							
				monocrotophos (0.030–0.089)	2							
				dimethoate (0.013–0.019)	0							
				pirimiphos-methyl (0.025)	0							
											cyfluthrin (0.055–0.429)	2
											cypermethrin (0.073–1.182)	4
											deltamethrin (0.424–1.030)	4
											fenvalerate (0.046–0.906)	2
Yard long bean	20	12	5	chlorpyrifos (0.002–0.033)	3	16	8	4	chlorpyrifos (0.003–0.018)	1		
				dicrotophos (0.104)	0				dicrotophos (0.001–0.060)	0		
				dimethoate (0.004–0.082)	1				ethion (0.001)	0		
				EPN (0.004–0.616)	4				profenophos (0.014)	0		
				malathion (0.002)	0							
				parathion-methyl (0.002)	0							
				profenophos (0.024–0.025)	0							
				triazophos (0.043)	0							
											cyfluthrin (0.096–0.619)	6
											cypermethrin (0.078–0.228)	1
				deltamethrin (0.434–0.681)	4							
				fenvalerate (0.092–0.553)	2							
				lamda-cyhalothrin (0.211–0.976)	4							
				permethrin (0.021–0.130)	0							
	4	3	NA	carbendazim (0.008–0.334)	1							

Supplement A. Continued.

Samples	Urban					Rural				
	N	No. of contaminated samples	No. of samples detected multi-residues	Detected pesticides; range or mean ^b (mg/kg)	No. exceeding MRL	N	No. of contaminated samples	No. of samples detected multi-residues	Detected OP pesticides; range or mean ^b (mg/kg)	No. exceeding MRL
Fruits	100	74 (74%)	38	chlorpyrifos diazinon dimethoate ethion fenitrothion malathion methamidophos methidathion monocrotophos parathion-methyl pirimiphos-ethyl pirimiphos-methyl profenophos prothiophos triazophos		201	170 (84%)	135	aldicarb carbaryl chlorpyrifos cyfluthrin cypermethrin deltamethrin diazinon dicrotophos dimethoate ethion fenvalerate imidacloprid abamectin 1a lamda-cyhalothrin malathion methamidophos monocrotophos permethrin profenophos prothiophos triazophos	
Apple	20	12	1	chlorpyrifos (0.003–0.161) ethion (0.001) fenitrothion (0.004–0.007)	1 0 0	19	7	5	chlorpyrifos (0.001–0.018) ethion (0.001) cyfluthrin (0.041–0.147) cypermethrin (0.010–0.290) deltamethrin (0.008–1.177) fenvalerate (0.014–0.196) lamda-cyhalothrin (0.024–0.385) permethrin (0.002–2.271)	2 0 2 1 8 0 10 5

Supplement A. Continued.

Samples	Urban					Rural				
	N	No. of contaminated samples	No. of samples detected multi-residues	Detected pesticides; range or mean ^b (mg/kg)	No. exceeding MRL	N	No. of contaminated samples	No. of samples detected multi-residues	Detected OP pesticides; range or mean ^b (mg/kg)	No. exceeding MRL
Asian pear	-	-	-	-	-	12	11	2	chlorpyrifos (0.001–0.038) dicotophos (0.001) ethion (0.001–0.003) cyfluthrin (0.013–0.513) cypermethrin (0.008–0.151) deltamethrin (0.023–0.397) fenvalerate (0.050–0.765) lamda-cyhalothrin (0.024–0.395) permethrin (0.011–0.340)	7 0 0 3 0 7 5 8 0
Dragon fruit	10	3	0	fenitrothion (0.004–0.011)	0	5	2	0	ethion (0.001–0.002)	0
Guava	20	20	8	chlorpyrifos (0.003–0.057) dimethoate (0.003–0.005) ethion (0.002–0.092) malathion (0.001–0.010) parathion-methyl (0.002–0.004) profenophos (0.156) prothiophos (0.007–0.0303)	8 0 0 0 0 1 0	12	9	5	chlorpyrifos (0.002–0.027) diazinon (0.003) dicotophos (0.001) ethion (0.001–0.224) methamidophos (0.006–0.008) prothiophos (0.005) cyfluthrin (0.011–2.79) cypermethrin (0.021–0.642) deltamethrin (0.105–0.540) fenvalerate (0.027–0.272) lamda-cyhalothrin (0.015–0.324) permethrin (0.015–0.492)	1 0 0 1 0 0 4 2 6 1 8 0
Grape	10	10	8	chlorpyrifos (0.003–0.038) dimethoate(0.002–0.120) ethion (0.001) malathion (0.005) methamidophos (0.025–0.030) methidathion (0.002) monocrotophos (0.008) pirimiphos-ethyl (0.001) pirimiphos-methyl (0.012) prothiophos (0.005–0.373)	2 3 0 0 2 0 0 0 0 0 2	2	2	2	chlorpyrifos (0.001–0.006) ethion (0.003) monocrotophos (0.001)	0 0 0

Supplement A. Continued.

Samples	Urban					Rural				
	N	No. of contaminated samples	No. of samples detected multi-residues	Detected pesticides; range or mean ^b (mg/kg)	No. exceeding MRL	N	No. of contaminated samples	No. of samples detected multi-residues	Detected OP pesticides; range or mean ^b (mg/kg)	No. exceeding MRL
Lychee	-	-	-	-	-	5	5	0	chlorpyrifos (0.001–0.002)	0
									ethion (0.001–0.498)	1
Mango	-	-	-	-	-	14	6	1	chlorpyrifos (0.001–0.022)	2
									ethion (0.001)	0
									triazophos (0.001)	0
									cyfluthrin (0.010–0.700)	3
									cypermethrin (0.006–0.155)	0
									deltamethrin (0.030–0.382)	6
									fenvalerate (0.003–1.024)	4
									lamda-cyhalothrin (0.030–0.186)	6
permethrin (0.007–0.491)	0									
Rose apple	-	-	-	-	-	10	6	4	diazinon (0.001)	0
									dicrotophos (0.001)	0
									dimethoate (0.005–0.007)	0
									ethion (0.001–0.096)	0
									cyfluthrin (0.020–0.131)	1
									cypermethrin (0.051–1.595)	3
									deltamethrin (0.020–2.739)	6
									fenvalerate (0.050–3.744)	3
									lamda-cyhalothrin (0.043–0.325)	6
									permethrin (0.019–0.200)	0
Strawberry	10	9	2	chlorpyrifos (0.003)	0	-	-	-	-	-
				diazinon (0.001)	0					
				ethion (0.002–0.467)	1					
				malathion (0.002)	0					
				traizophos (0.001–0.490)	1					

Supplement A. Continued.

Samples	Urban					Rural						
	N	No. of contaminated samples	No. of samples detected multi-residues	Detected pesticides; range or mean ^b (mg/kg)	No. exceeding MRL	N	No. of contaminated samples	No. of samples detected multi-residues	Detected OP pesticides; range or mean ^b (mg/kg)	No. exceeding MRL		
Tangerine	20	19	19	chlorpyrifos (0.003–0.315)	9	22	22	20	chlorpyrifos (0.002–0.150)	16		
				diazinon (0.001)	0				diazinon (0.001–0.014)	0		
				dimethoate (0.017–0.131)	1				dicrotophos (0.121)	0		
				ethion (0.002–0.467)	1				dimethoate (0.005–0.157)	2		
				malathion (0.002–0.068)	1				ethion (0.002–1.130)	11		
				methamidophos (0.023–0.069)	8				malathion (0.001–0.288)	2		
				pirimiphos-methyl (0.005–0.019)	0				methamidophos (0.009–0.052)	12		
				profenophos (0.008–0.303)	0				monocrotophos (0.063)	1		
				triazophos (0.001–0.490)	1				prothiophos (0.001–0.002)	0		
									profenophos (0.017–1.196)	10		
									triazophos (0.151–0.325)	2		
									cyfluthrin (0.031–0.272)	6		
									cypermethrin (0.106–5.398)	12		
									deltamethrin (0.228–0.788)	10		
									fenvalerate (0.112–0.942)	4		
									lamda-cyhalothrin (0.026–0.971)	12		
									permethrin (0.012–0.172)	0		
											chlorpyrifos (0.002–0.211)	0
											dimethoate (0.018–0.629)	0
				ethion (0.003–2.887)	21							
				malathion (0.002–0.360)	0							
				methamidophos (0.001–0.118)	0							
				profenophos (0.002–0.211)	2							
				cyfluthrin (0.15–86.1)	25							
				cypermethrin (0.01–3.95)	1							
				fenvalerate (0.01–0.47)	18							
				lamda-cyhalothrin (0.01–0.80)	6							
				permethrin (0.01–0.80)	8							
				aldicarb (0.45)	1							
				carbaryl (0.14–0.31)	0							
				imidacloprid (0.003–0.048)	0							
				abamectin 1a (0.003–0.035)	13							
Watermelon	10	1	0	chlorpyrifos (0.032)	1	-	-	-	-	-		

Supplement B. Multiple pesticide residues detected in samples collected from urban and rural markets in upper northern Thailand during 2007-2013.

Pesticide	MRL* (mg/kg)	Urban							Rural						
		N	N detected	% detected	exceeded	% exceeded	Mean mg/kg	Min-Max mg/kg	N	N detected	% detected	exceeded	% exceeded	Mean mg/kg	Min-Max mg/kg
Organophosphate															
Azinphos-ethyl	0.05-1 Co	300	ND	ND	ND	ND	ND	ND	337	ND	ND	ND	ND	ND	ND
Azinphos-methyl	0.5-1 Eu	300	1	0.3	0	0.0	0.088	0.088-0.088	337	ND	ND	ND	ND	ND	ND
Chlopyrifos	0.01-2 Co	300	107	35.7	61	20.3	0.076	0.002-1.394	337	92	40.5	43	18.9	0.052	0.001-1.596
Diazinon	0.05-0.5 Co	300	10	3.3	4	1.3	0.094	0.0005-0.402	337	12	5.3	0	0.0	0.004	0.002-0.014
Dicrotophos	0.5 Ta	300	21	7.0	0	0.0	0.056	0.003-0.403	337	37	16.3	3	1.3	0.095	0.001-0.973
Dimethoate	0.05-1 Co	300	28	9.3	7	2.3	0.032	0.001-0.131	337	11	4.8	3	1.3	0.041	0.001-0.159
EPN	0.02-0.1 Ja	300	8	2.7	4	1.3	0.151	0.004-0.616	337	ND	ND	ND	ND	ND	ND
Ethion	0.1-2 Eu	300	41	13.7	10	3.3	0.144	0.001-1.658	337	76	33.5	14	6.2	0.101	0.0002-1.279
Fenitrothion	0.5 Co	300	8	2.7	0	0.0	0.027	0.004-0.161	337	ND	ND	ND	ND	ND	ND
Malathion	0.2-8 Co	300	17	5.7	2	0.7	0.072	0.002-0.776	337	9	4.0	1	0.4	0.083	0.001-0.288
Methamidophos	0.01-0.3 Ja	300	20	6.7	17	5.7	0.069	0.005-0.639	337	21	9.3	11	4.8	0.017	0.002-0.052
Methidathion	0.05-1 Co	300	2	0.7	0	0.0	0.002	0.002-0.002	337	ND	ND	ND	ND	ND	ND
Mevinphos	0.05-0.5 Ja	300	2	0.7	0	0.0	0.020	0.011-0.030	337	ND	ND	ND	ND	ND	ND
Monocrotophos	0.05-1 Ja	300	6	2.0	2	0.7	0.031	0.003-0.089	337	4	1.8	3	1.3	0.095	0.001-0.313
Parathion-methyl	0.05-0.5 Co	300	3	1.0	0	0.0	0.002	0.001-0.004	337	2	0.9	0	0.0	0.011	0.002-0.020
Pirimiphos-ethyl	NA	300	8	2.7	0	0.0	0.011	0.001-0.022	337	2	0.9	0	0.0	0.001	0.001-0.001
Pirimiphos-methyl	0.05-5 Ja	300	15	5.0	0	0.0	0.011	0.002-0.034	337	ND	ND	ND	ND	ND	ND
Profenophos	0.05-0.2 Ja	300	23	7.7	14	4.7	0.097	0.008-0.303	337	26	11.5	14	6.2	0.152	0.001-1.196
Prothiofos	0.2-2 Ja	300	11	3.7	2	0.7	0.068	0.005-0.373	337	5	2.2	0	0.0	0.003	0.001-0.005
Triazophos	0.1 Co	300	5	1.7	1	0.3	0.110	0.001-0.490	337	5	2.2	2	0.9	0.097	0.001-0.325
Carbamates															
3-Hydroxy carbofuran	NA	-	-	-	-	-	-	-	100	ND	ND	ND	ND	ND	ND
Aldicarb	0.3 Ja	-	-	-	-	-	-	-	100	1	1.1	1	1.1	0.45	0.45
Aldicarb sulfone	NA	-	-	-	-	-	-	-	100	ND	ND	ND	ND	ND	ND
Aldicarb sulfoxide	NA	-	-	-	-	-	-	-	100	ND	ND	ND	ND	ND	ND
Baygon	NA	-	-	-	-	-	-	-	100	ND	ND	ND	ND	ND	ND
Carbaryl	1.5 Th	-	-	-	-	-	-	-	100	2	2.2	2	0	NA	0.14-0.31
Carbofuran	0.5 Ko	-	-	-	-	-	-	-	100	ND	ND	ND	ND	ND	ND
Methiocarb	NA	-	-	-	-	-	-	-	100	ND	ND	ND	ND	ND	ND
Methomyl	1 Th	-	-	-	-	-	-	-	100	ND	ND	ND	ND	ND	ND
Oxamyl	5 Ko	-	-	-	-	-	-	-	100	ND	ND	ND	ND	ND	ND

Supplement B. Continued.

Pesticide	MRL* (mg/kg)	Urban							Rural						
		N	N detected	% detected	exceeded	% exceeded	Mean mg/kg	Min-Max mg/kg	N	N detected	% detected	exceeded	% exceeded	Mean mg/kg	Min-Max mg/kg
Pyrethroids															
Cyfluthrin	0.1–2 Co	-	-	-	-	-	-	-	171	106	62.0	59	34.5	0.19	0.01–2.79
Cypermethrin	0.2–2 Co	-	-	-	-	-	-	-	171	135	79.0	37	21.6	0.42	0.01–8.39
Deltamethrin	0.02–2 Co	-	-	-	-	-	-	-	171	99	57.9	97	56.7	0.92	0.008–17.45
Fenvalerate	0.2–2 Co	-	-	-	-	-	-	-	171	104	61.0	51	29.8	0.42	0.0025–3.74
Lambda-cyhalothrin	0.05–0.5 Co	-	-	-	-	-	-	-	171	127	74.3	112	65.5	0.28	0.02–1.07
Permethrin	0.5–5 Co	-	-	-	-	-	-	-	171	114	66.7	5	2.9	0.14	0.002–2.27
Other insecticides															
Abamectin 1a	0.01 Th	-	-	-	-	-	-	-	100	40	44.0	13	13	NA	0.003–0.035
Abamectin 1b	0.01 Th	-	-	-	-	-	-	-	100	ND	ND	ND	ND	ND	ND
Imidacloprid	0.5 Ko	-	-	-	-	-	-	-	100	79	86.8	0	0	NA	0.003–0.048
Fungicides															
Carbendazim	0.05–0.5 Co (7 K)	32	21	65.6	5	15.6	0.047	0.008–0.334	100	27	29.7	0	0	NA	0.006–0.077
Dithiocarbamates (Mancozeb)	2.0 Th	54	10	18.5	3	5.6	4.18	0.120–30.64	-	-	-	-	-	-	-